

Muon anomalous magnetic moment in a Calculable Model with one Extra Dimension

Giacomo Cacciapaglia^{a,1}, Marco Cirelli^{a,2}, Giampaolo Cristadoro^{b,3}

^a *Scuola Normale Superiore and INFN, Piazza dei Cavalieri 7, I-56126 Pisa, Italy*

^b *‘Enrico Fermi’ Dep. of Physics, University of Pisa, via Buonarroti 2, I-56126 Pisa, Italy*

Abstract

In the framework of a recently proposed extension of the Standard Model, with $N = 1$ Super-Symmetry in 5 dimensions, compactified on $\mathbb{R}^1/\mathbb{Z}_2 \times \mathbb{Z}'_2$, we compute the muon anomalous magnetic moment at one loop to order $(M_W R)^2$, where R is the compactification radius. We find the corrections to be small with respect to the SM pure weak contribution and not capable of explaining the present discrepancy between theory and experiment for any sensible value of R .

PACS: 11.10.Kk, 12.60.Jv, 13.40.Em

¹cacciapa@cibs.sns.it

²mcirelli@cibs.sns.it

³cristado@cibs.sns.it

1 Introduction

Models with extra (space) dimensions often fail to produce quantitative predictions for physical observables and, as a consequence, can hardly be ruled out or confirmed by present energy experiments. On the other hand, in [1] a model was proposed where calculability is achieved for a number of physical quantities. In fact, in spite of the non renormalizability of 5 dimensional Yang-Mills theory, they are finite and cut-off independent, owing to the underlying supersymmetric structure. The insensitivity to ultraviolet physics of such observables is one of the main goals of the model. The branching ratio of $B \rightarrow X_s \gamma$ [2] and the Higgs production via gluon fusion [3] have already been computed, while in the present paper we deal with the muon anomalous magnetic moment a_μ .

The measurement of a_μ is one of the most stringent tests for “new physics” scenarios, thanks to its current impressive precision. Moreover, the recent results of E821 experiment at BNL [4] have highlighted a discrepancy with the present Standard Model prediction and have therefore made even more interesting the comparison with reliable predictions made by new models. In particular, attempts have been made in order to study the effect of extra space dimensions [5].

Let us briefly review the present status of the SM prediction [6]. The QED contribution has been computed up to order α^5 and it gives the core of the experimental value: $a_\mu^{QED} = (11\,658\,470.0 \pm 0.3) \cdot 10^{-10}$. The purely weak terms are also known with a small uncertainty, up to two loops: $a_\mu^{weak} = (15.1 \pm 0.4) \cdot 10^{-10}$. On the contrary, hadronic loop corrections provide relevant contributions but also carry the greatest theoretical uncertainty [7]. In fact, since it is quite hard to deal with low energy QCD, experimental inputs are also needed. Light-by-light scattering yields $a_\mu^{lbl} = -(8.6 \pm 3.2) \cdot 10^{-10}$ and hadronic vacuum polarization $a_\mu^{hvp} = (684.9 \pm 6.4) \cdot 10^{-10}$. Summing up all these contributions and uncertainties, the SM prediction is:

$$a_\mu^{SM} = (11\,659\,162.0 \pm 7.5) \cdot 10^{-10}$$

On the other hand, the present world average for the experimental value (including BNL results) is [6]:

$$a_\mu^{exp} = (11\,659\,202.3 \pm 15.1) \cdot 10^{-10}$$

This leads to a discrepancy a bit larger than 2σ :

$$a_\mu^{exp} - a_\mu^{SM} = (40.3 \pm 16.9) \cdot 10^{-10} \tag{1}$$

Actually, this result is not well established. On the theoretical side, recent results [8] [9] claim that the light-by-light contribution has opposite sign, thus reducing the discrepancy to about 1σ . At the same time, the error on the experimental value should be improved by further results at BNL.

In this work we compute the corrections Δa_μ to the anomalous magnetic moment due to the new fields in the theory, up to order $(M_W R)^2$ where R is the compactification radius of the 5th dimension.

2 The model

In this section we briefly summarize the main features of the model [1] we are working with. It is a 5D theory with $N=1$ supersymmetry, compactified on $\mathbb{R}^1/\mathbb{Z}_2 \times \mathbb{Z}'_2$, and the gauge group is the Standard Model one: $SU(3) \times SU(2) \times U(1)$. The fields content is made up of the vector hypermultiplets and a matter hypermultiplet for each SM matter field Q, U, D, L, E, H , all living in the bulk.

From a 4D point of view, a matter hypermultiplet splits into a pair of chiral supermultiplets with conjugate quantum numbers (property that we indicate with a c), while a vector hypermultiplet splits into a vector and a chiral supermultiplet. Under the orbifolding, global supersymmetry is completely broken *à la* Scherk-Schwarz, leaving the SM fields as the only zero modes. Anyhow, restricted local supersymmetric transformations still hold. The most general Lagrangian, according to this set of symmetries (gauge group, orbifold parities and local supersymmetry), is:

$$\mathcal{L}(x, y) = \mathcal{L}_5 + \delta(y) \mathcal{L}_4 + \delta(y - \pi R/2) \mathcal{L}'_4 \quad (2)$$

where $\mathcal{L}_5$ is $N = 1$ supersymmetric in 5D whereas $\mathcal{L}_4$ and $\mathcal{L}'_4$ are 4D lagrangians invariant under different $N = 1$ supersymmetries valid on the fixed points of the two parities. To be consistent with ref. [1], we will write all hypermultiplets in terms of supermultiplets of the 4-dimensional $N=1$ supersymmetry located on the brane $y = 0$. So, the gauge vector and matter hypermultiplets are:

$$(W_\rho, \lambda) \oplus (\lambda', \Sigma = \frac{1}{\sqrt{2}}(\sigma + iW_5))$$

$$(\psi_X, \phi_X) \oplus (\psi_X^c, \phi_X^c)$$

Under the two parities all the fields have definite transformation properties, given in Table 1 together with the corresponding eigenfunctions and the spectrum of every tower of Kaluza-Klein states.

A_μ, ψ_M, ϕ_H	$(+, +)$	$\cos \frac{2n}{R}y$	$\frac{2n}{R}, n = 0, 1, 2, 3...$
λ, ϕ_M, ψ_H	$(+, -)$	$\cos \frac{2n-1}{R}y$	$\frac{2n-1}{R}, n = 1, 2, 3...$
$\psi_\Sigma, \phi_M^c, \psi_H^c$	$(-, +)$	$\sin \frac{2n-1}{R}y$	$\frac{2n-1}{R}, n = 1, 2, 3...$
$\phi_\Sigma, \psi_M^c, \phi_H^c$	$(-, -)$	$\sin \frac{2n}{R}y$	$\frac{2n}{R}, n = 1, 2, 3...$

Table 1: Gauge, Matter and Higgs fields content of the theory with their orbifolding properties

In [10] it is shown that in the model under consideration the hypercharge current has an anomaly, which is localized on the two branes. A relevant feature is that, from a 4 dimensional point of view, the integrated anomaly vanishes. Gauge invariance can however be recovered with a suitable modification of the theory. The overall consistency of the model modified by the addition of a Chern-Simons term, for instance, is currently under examination. Such modifications, however, should not affect the calculation

presented in this paper, since anomalous contributions only enter at higher order.

The terms in the 5D action, that are relevant for the following calculation, are:

$$\begin{aligned}
S = \int d^5x \quad & \left\{ -\frac{1}{4} \mathbf{Tr} F_{MN} F^{MN} - \mathbf{Tr} \left[i\lambda \sigma^\rho \mathcal{D}_\rho \bar{\lambda} - i\lambda' \sigma^\rho \mathcal{D}_\rho \bar{\lambda}' \right] + i\psi_X \sigma^\rho \mathcal{D}_\rho \bar{\psi}_X + i\psi_X^c \sigma^\rho \mathcal{D}_\rho \bar{\psi}_X^c + \right. \\
& + (\mathcal{D}_M \phi_X)^\dagger (\mathcal{D}^M \phi_X) + (\mathcal{D}_M \phi_X^c)^\dagger (\mathcal{D}^M \phi_X^c) + \sqrt{2} g_{(5)} \left[\phi_X^\dagger \lambda \psi_X - \psi_X^c \lambda \phi_X^{c\dagger} + h.c. \right] + \\
& + \sqrt{2} g_{(5)} \left[\psi_X^c \lambda' \phi_X + \phi_X^c \lambda' \psi_X + \psi_X^c \Sigma \psi_X + h.c. \right] + \\
& + \delta(y - \frac{\pi}{2} R) \lambda_\mu \left[\psi_L \psi_E \phi_{H^0}^\dagger + \psi_L \psi_{H^0}^c \phi_E^\dagger + \psi_E \psi_{H^0}^c \phi_L^\dagger + \right. \\
& \left. \left. - (\psi_\nu \psi_E \phi_{H^+}^\dagger + \psi_\nu \psi_{H^+}^c \phi_E^\dagger + \psi_E \psi_{H^+}^c \phi_\nu^\dagger) \right] \right\} \quad (3)
\end{aligned}$$

where the fermionic fields are Weyl spinors, the gauge fields W_M , λ , λ' and σ contains the group generators, e.g. $\lambda \equiv \lambda^a T^a$ (the generators being normalized according to $\mathbf{Tr}(T^a T^b) = \delta^{ab}$) and the index X can run over E , L , H . The gauge coupling g is intended to be the $U_Y(1)$ coupling g' or the $SU_L(2)$ coupling g according to the gauge fields involved.

From now on, for the four gauge hypermultiplets we use the notation:

$$\begin{aligned}
QED & \quad (A_\rho, \tilde{\gamma}, \tilde{\gamma}_c, \Sigma_\gamma) \\
Z's & \quad (Z_\rho, \tilde{z}, \tilde{z}_c, \Sigma_z) \\
W's & \quad (W_\rho^\pm, \tilde{w}^\pm, \tilde{w}_c^\pm, \Sigma_w^\pm)
\end{aligned}$$

(we remind that, also in this case, the subscribed c only means that the second spinors have conjugate quantum numbers).

2.1 Mass Eigenstates

The brane interactions introduce mass mixing among the states of muon and smuon KK towers: since it will be useful for our calculation to work with the mass eigenstates, we proceed to determine eigenvalues and eigenvectors from the very beginning (see e.g. the discussion in [2]). For the fermions, the mass eigenvalues turn out to be:

$$m_0 = m \equiv \frac{\epsilon}{R}, \quad m_n^\pm = m \pm \frac{2n}{R} \quad (n = 1, 2, \dots)$$

corresponding to mass eigenvectors whose Weyl spinor components we denote by (μ_0, μ_0^c) and $(\mu_n^\pm, \mu_n^{c\pm})$. Thus m is the mass of the ordinary muon. For the scalars:

$$\tilde{m}_n^\pm = \frac{(2n-1)}{R} \pm m \quad (n = 1, 2, \dots)$$

corresponding to eigenvectors $\Phi_n^\pm$ and $\Phi_n^{c\pm}$.

The interaction eigenstates¹ are expressed in terms of the mass eigenstates (at order ϵ) by

$$\psi_{L,0} = \mu_0 + \epsilon \sum_n \frac{1}{2n} (\mu_n^+ - \mu_n^-) \quad (4)$$

$$\psi_{L,l} = \frac{1}{\sqrt{2}} \left\{ \left(1 + \frac{\epsilon}{2(2n)}\right) \mu_l^+ + \left(1 - \frac{\epsilon}{2(2n)}\right) \mu_l^- + \epsilon \sum_n \kappa_{nl} (\mu_n^+ - \mu_n^-) \right\} \quad (5)$$

$$\psi_{E,l}^c = \frac{1}{\sqrt{2}} \left\{ \left(1 - \frac{\epsilon}{2(2n)}\right) \mu_l^+ - \left(1 + \frac{\epsilon}{2(2n)}\right) \mu_l^- - \epsilon \sum_n \kappa_{ln} (\mu_n^+ + \mu_n^-) \right\} - \frac{\sqrt{2}}{2l} \epsilon \mu_0 \quad (6)$$

$$\phi_{L,l} = \frac{1}{\sqrt{2}} \left\{ \left(1 + \frac{\epsilon}{2(2n-1)}\right) \Phi_l^+ + \left(1 - \frac{\epsilon}{2(2n-1)}\right) \Phi_l^- + \epsilon \sum_n J_{ln} (\Phi_n^+ - \Phi_n^-) \right\} \quad (7)$$

$$\phi_{E,l}^c = \frac{1}{\sqrt{2}} \left\{ - \left(1 - \frac{\epsilon}{2(2n-1)}\right) \Phi_l^+ + \left(1 + \frac{\epsilon}{2(2n-1)}\right) \Phi_l^- + \epsilon \sum_n J_{nl} (\Phi_n^+ + \Phi_n^-) \right\} \quad (8)$$

with $\kappa_{ln} = \frac{2(2l)}{(2l)^2 - (2n)^2}$ and $J_{ln} = \frac{2(2l-1)}{(2l-1)^2 - (2n-1)^2}$, and zero for $n = l$. Expressions which are identical to (4), (5), (6) hold for $(\psi_{E,0}, \psi_{E,n}, \psi_{L,n}^c)$ if $(\mu_0, \mu_n^\pm)$ are replaced by $(\mu_0^c, \mu_n^{c\pm})$; for (ϕ_L^c, ϕ_E) it is sufficient to replace $\Phi_n^\pm$ with $\Phi_n^{c\pm}$ in equations (7), (8).

2.2 Couplings to the mass eigenstates

Starting from eq. (3), one has to integrate out the 5^{th} dimension y and rewrite the interaction vertices in terms of the mass eigenstates (4)-(8). Taking into account momentum conservation along the 5^{th} direction, p_5 , the bulk interactions give rise to several trilinear vertices. The muon masses and Higgs vertices, on the other hand, being located on the brane, violate p_5 -conservation and are proportional to ϵ . As we are interested in $\mathcal{O}(R^2)$ contributions, only graphs where the internal lines carry the same index n , in terms of the mass eigenstates, have to be taken into account. This simplifies the calculation of the relevant vertices. As an explicit example, the interaction terms between KK photons and the various components in the muon towers, in the mass eigenstate basis, are:

$$\begin{aligned} e \bar{\psi}_{L,0} \bar{\sigma}_\rho \psi_{L,n} A_n^\rho &= e \bar{\mu}_0 \bar{\sigma}_\rho \frac{1}{\sqrt{2}} \left\{ \left(1 - \frac{\epsilon}{2(2n)}\right) \mu_n^+ + \left(1 + \frac{\epsilon}{2(2n)}\right) \mu_n^- \right\} A_n^\rho \\ \frac{e}{\sqrt{2}} \bar{\psi}_{E,2n}^c \bar{\sigma}_\rho \psi_{E,n}^c A_n^\rho &= -e \frac{\epsilon}{2(2n)} \bar{\mu}_0 \bar{\sigma}_\rho \frac{1}{\sqrt{2}} \{ \mu_n^+ - \mu_n^- \} A_n^\rho \end{aligned}$$

In this particular case, summing up the two contributions, the couplings of order ϵ vanish, so that the net result is:

¹Note that in this paper the brane lagrangians $\mathcal{L}_4$ and $\mathcal{L}'_4$ are reversed respect to ref [1]. For more details, see appendix A in [3].

$$e\bar{\mu}_0\bar{\sigma}_\rho\frac{1}{\sqrt{2}}(\mu_n^+ + \mu_n^-)A_n^\rho \quad (9)$$

This allows to recognize $e/\sqrt{2}$ as the correct coupling with a KK photon.

On the other hand, the interactions of the zero mode gauge bosons are still diagonal in n and have a universal coupling, e.g. $e = e_{(5)}/\sqrt{2\pi R}$ for the photon. Note that this relation holds for all gauge couplings, so in the following $g = g_{(5)}/\sqrt{2\pi R}$ is the 4D gauge coupling, while $s = \sin\theta_W$ and $c = \cos\theta_W$ are functions of the usual SM Weinberg angle.

3 Calculation of a_μ

The effective Lagrangian term which fixes the notation for the anomalous magnetic moment $a_\mu = (g_\mu - 2)/2$ is:

$$\mathcal{L} = \frac{ie}{2m} a_\mu (\bar{\mu}\sigma_{\rho\sigma}F^{\rho\sigma}\mu) \quad (10)$$

In the Feynman-'t Hooft gauge ($\xi = 1$, see below), one can identify the five general types of diagrams in fig. 1, each giving a contribution Δa_μ whose explicit expressions are listed in app. A.




Figure 1: General patterns of graphs contributing to the $(g - 2)$.

As in the following we will distinguish between diagrams with chirality flip on the external or internal fermionic line, in eq. (51)-(54) it is easy to trace back the external cross contribution and the internal

$$\mathcal{L}_{mass} = -\partial_\rho A^\rho \left(\frac{2n}{R} A_5 \right) - \frac{1}{2} \left(\frac{2n}{R} \right)^2 \sigma_\gamma^2 - \frac{1}{2\xi} \left(\partial_\rho A^\rho - \xi \frac{2n}{R} A_5 \right)^2 \quad (14)$$

where the R_ξ gauge fixing term is there to cancel the mixing terms. It is convenient to work in the $\xi = 1$ gauge: we obtain a photon propagator with no longitudinal modes and two scalars σ_γ and A_5 (with equal mass) that can be combined in the complex scalar Σ_γ . The interaction terms are then:

$$\sqrt{2}e \left(\psi_E^c \Sigma_\gamma \psi_E + \psi_L^c \Sigma_\gamma \psi_L + h.c. \right) \quad (15)$$

so that at the relevant order in ϵ , only an external cross graph is possible. Equivalently, in terms of the real scalar fields it happens that the internal cross diagram with an A_5 field is cancelled out by the same diagram with a σ_γ . The external cross diagram is of type *II* so that the resulting Δa_μ is

$$\mu_0^c \text{---}\times\text{---}\overset{\mu^\pm}{\text{---}}\mu_0 \quad \Rightarrow \quad \frac{e^2}{16\pi^2} \frac{\epsilon^2}{(2n)^2} \int dx [4x(1-x)^2] \quad (16)$$

Σ_γ

3.1.3 Fermions

The gauge fermions $\tilde{\gamma}$ and $\tilde{\gamma}_c$ are mixed by the kinetic term along the 5th direction. They are involved in the following interaction terms

$$\begin{aligned} & -\sqrt{2}e\mu_0\tilde{\gamma}_n \frac{1}{\sqrt{2}} \left\{ \left(\Phi_n^{\dagger-} - \Phi_n^{\dagger+} \right) + \frac{\epsilon}{2n-1} \left(\Phi_n^{\dagger-} + \Phi_n^{\dagger+} \right) \right\} \\ & +\sqrt{2}e\mu_0^c\tilde{\gamma}_{c,n} \frac{1}{\sqrt{2}} \left\{ \left(\Phi_n^+ + \Phi_n^- \right) + \frac{\epsilon}{2n-1} \left(\Phi_n^+ - \Phi_n^- \right) \right\} \end{aligned} \quad (17)$$

and in similar expressions (replacing Φ with Φ^c) for the interactions of μ_0^c with $\tilde{\gamma}_n$ and μ_0 with $\tilde{\gamma}_{c,n}$. From graph *II*, one gets

$$\left. \begin{array}{l} \mu_0^c \text{---}\times\text{---}\overset{\Phi^\pm}{\text{---}}\mu_0 \\ \quad \quad \quad \tilde{\gamma} \\ \mu_0^c \text{---}\times\text{---}\overset{\Phi^{c\pm}}{\text{---}}\mu_0 \\ \quad \quad \quad \tilde{\gamma}_c \end{array} \right\} \Rightarrow -\frac{e^2}{16\pi^2} \frac{\epsilon^2}{(2n-1)^2} \int dx [4x(1-x)^2] \times 2 \quad (18)$$

$$\left. \begin{array}{c} \text{Diagram 1: } \mu_0^c \text{ --- } \mu_0 \text{ with loop } \tilde{\gamma}^c, \tilde{\gamma} \text{ and boson } \Phi^\pm \\ \text{Diagram 2: } \mu_0^c \text{ --- } \mu_0 \text{ with loop } \tilde{\gamma}, \tilde{\gamma}^c \text{ and boson } \Phi^{c\pm} \end{array} \right\} \Rightarrow \frac{e^2}{16\pi^2} \frac{\epsilon^2}{(2n-1)^2} \int dx [8x(1-x)^2] \times 2 \quad (19)$$

3.1.4 Total QED contribution

First of all, in order to test the computation, we consider the supersymmetric limit in which both fermions and bosons have the same parities and therefore the same bulk mass. In this case, we find that the total contributions of the KK modes vanishes, as expected. Of course, the appearance of zero modes for the matter and gauge partners also cancels the SM result.

Then, we collect all the QED contributions and we obtain:

$$\Delta a_\mu^{QED} = \frac{e^2}{16\pi^2} (mR)^2 \frac{\pi^2}{18} \quad (20)$$

3.2 Neutral Weak sector

In this section we consider all the diagrams involving Z hypermultiplet fields and neutral Higgs. We define

$$\begin{aligned} g_L &= \frac{1}{c} \left(-\frac{1}{2} + s^2 \right) g \\ g_R &= -\frac{s^2}{c} g \end{aligned}$$

The one loop SM result [12] comes from graph *IV* with a zero mode Z boson and couplings $g_- = g_L$, $g_+ = -g_R$:

$$a_\mu^Z = \frac{g^2}{48\pi^2} \frac{m^2}{M_Z^2} \frac{4s^4 - 2s^2 - 1}{c^2} \quad (21)$$

The discussion for the KK modes mimics the QED case, thus in the following we only point out the differences.

3.2.1 KK vectors

The interaction lagrangian consists in

$$\bar{\mu}_0 \bar{\sigma}_\rho \frac{1}{\sqrt{2}} \{j_L^+ \mu_n^+ + j_L^- \mu_n^-\} Z_n^\rho + \bar{\mu}_0^c \bar{\sigma}_\rho \frac{1}{\sqrt{2}} \{j_R^+ \mu_n^{c+} + j_R^- \mu_n^{c-}\} Z_n^\rho + h.c. \quad (22)$$

where:

$$\begin{aligned} j_L^+ &= g_L + \frac{\epsilon}{2(2n)} (g_L + g_R) & j_R^+ &= -g_R - \frac{\epsilon}{2(2n)} (g_R + g_L) \\ j_L^- &= g_L - \frac{\epsilon}{2(2n)} (g_L + g_R) & j_R^- &= -g_R + \frac{\epsilon}{2(2n)} (g_R + g_L) \end{aligned} \quad (23)$$

The two Δa_μ then amount to

$$\mu_0^c \text{---} \times \text{---} \mu^\pm \text{---} \mu_0 \quad \Rightarrow \quad \frac{1}{16\pi^2} \frac{\epsilon^2}{(2n)^2} \int dx (g_L^2 + g_R^2) [2x(1-x)^2 - 4x(1-x)] \quad (24)$$

Z_ρ

$$\begin{aligned} \mu_0^c \text{---} \mu^{c\pm} \text{---} \mu^\pm \text{---} \mu_0 \quad \Rightarrow \quad & -\frac{1}{16\pi^2} \frac{\epsilon^2}{(2n)^2} \int dx \left[\left(g_L g_R + \frac{1}{2} (g_L + g_R)^2 \right) 8x(1-x) \right. \\ & \left. - g_L g_R 16x(1-x)^2 \right] \end{aligned} \quad (25)$$

Z_ρ

3.2.2 Scalars

With respect to the QED case (14), the lagrangian for scalar fields is a bit more complicated, involving also mixing terms with φ_{H^0} , the imaginary part of ϕ_{H^0} . With the gauge fixing piece already included, the lagrangian for each mode n is:

$$\begin{aligned} \mathcal{L}_{mass} = & -\partial_\rho Z^\rho \left(\frac{2n}{R} Z_5 - M_Z \varphi_{H^0} \right) - \frac{1}{2} \left[\frac{2n}{R} \varphi_{H^0} - M_Z Z_5 \right]^2 - \frac{1}{4} M_Z^2 \varphi_{H^0}^2 - \frac{1}{2} \left(\frac{2n}{R} \right)^2 \sigma_Z^2 \\ & - \frac{1}{2\xi} \left[\partial_\rho Z^\rho - \xi \left(\frac{2n}{R} Z_5 - M_Z \varphi_{H^0} \right) \right]^2 \end{aligned} \quad (26)$$

Vector-scalar mixings are again cancelled and, for $\xi = 1$, there are no off-diagonal terms $\varphi_{H^0} Z_5$, so the three real scalars σ_Z , Z_5 , φ_{H^0} are the mass eigenstates. The first two combine in the complex field Σ_Z in terms of which, at the relevant order in ϵ , only the external cross graph is possible:

$$\mu_0^c \text{---}\times\text{---}\overset{\mu^\pm}{\text{---}}\mu_0 \Rightarrow \frac{1}{16\pi^2} \frac{\epsilon^2}{(2n)^2} \int dx (g_L^2 + g_R^2) 2x(1-x)^2 \quad (27)$$

3.2.3 Fermions

The mixing matrix for the four neutralinos $\tilde{z}$, $\tilde{z}_c$, ψ_{H^0} , $\psi_{H^0}^c$, for each mode n , is:

$$\mathcal{L}_{mass} = (\psi_{H^0}, \tilde{z}) \begin{pmatrix} \frac{2n-1}{R} & M_Z \\ -M_Z & \frac{2n-1}{R} \end{pmatrix} \begin{pmatrix} \psi_{H^0}^c \\ \tilde{z}_c \end{pmatrix} \quad (28)$$

They are involved in the following interaction terms:

$$\begin{aligned}
& \sqrt{2}\mu_0\tilde{z}_n\frac{1}{\sqrt{2}}\left\{g_L(\Phi_n^{\dagger-}-\Phi_n^{\dagger+})+(g_L-g_R)\frac{\epsilon}{2(2n-1)}(\Phi_n^{\dagger-}+\Phi_n^{\dagger+})\right\}+ \\
& +\sqrt{2}\mu_0^c\tilde{z}_n\frac{1}{\sqrt{2}}\left\{g_L\leftrightarrow g_R\quad\text{and}\quad\Phi\leftrightarrow\Phi^c\right\}+ \\
& +\sqrt{2}\mu_0^c\tilde{z}_n^c\frac{1}{\sqrt{2}}\left\{g_R(\Phi_n^-+\Phi_n^+)-(g_R-g_L)\frac{\epsilon}{2(2n-1)}(\Phi_n^--\Phi_n^+)\right\}+ \\
& +\sqrt{2}\mu_0\tilde{z}_n^c\frac{1}{\sqrt{2}}\left\{g_L\leftrightarrow g_R\quad\text{and}\quad\Phi\leftrightarrow\Phi^c\right\}+ \\
& +2\frac{\epsilon}{v}\left\{\mu_0\frac{1}{\sqrt{2}}(\Phi_n^-+\Phi_n^+)+\mu_0^c\frac{1}{\sqrt{2}}(\Phi_n^{c-}+\Phi_n^{c+})\right\}\psi_{H,n}^c
\end{aligned} \tag{29}$$

The external cross graphs are readily computed:

$$\left. \begin{array}{c} \mu_0^c \text{---} \times \text{---} \overbrace{\text{---} \text{---}}^{\Phi^\pm} \text{---} \mu_0 \\ \quad \quad \quad \underbrace{\hspace{1.5cm}}_{\tilde{z}} \\ \mu_0^c \text{---} \times \text{---} \overbrace{\text{---} \text{---}}^{\Phi^{c\pm}} \text{---} \mu_0 \\ \quad \quad \quad \underbrace{\hspace{1.5cm}}_{\tilde{z}_c} \end{array} \right\} \Rightarrow -\frac{1}{16\pi^2} \frac{\epsilon^2}{(2n-1)^2} \int dx (g_L^2 + g_R^2) 2x(1-x)^2 \times 2 \quad (30)$$

and it is possible to write three internal cross diagrams, using also the off-diagonal propagator in eq. (28):

$$\begin{array}{c} \mu_0^c \text{---} \text{---} \text{---} \mu_0 \\ \text{---} \Phi^\pm \text{---} \\ \text{---} \tilde{z}^c \text{---} \times \text{---} \tilde{z} \end{array} \Rightarrow -\frac{1}{16\pi^2} \frac{\epsilon^2}{(2n-1)^2} \int dx [(g_L^2 - g_R^2)2x(1-x) + g_L g_R 8x(1-x)^2] \quad (31)$$

$$\begin{array}{c} \mu_0^c \text{---} \text{---} \text{---} \mu_0 \\ \text{---} \Phi^{c\pm} \text{---} \\ \text{---} \tilde{z} \text{---} \times \text{---} \tilde{z}^c \end{array} \Rightarrow -\frac{1}{16\pi^2} \frac{\epsilon^2}{(2n-1)^2} \int dx [(g_R^2 - g_L^2)2x(1-x) + g_L g_R 8x(1-x)^2] \quad (32)$$

$$\begin{array}{c} \mu_0^c \text{---} \text{---} \text{---} \mu_0 \\ \text{---} \Phi^{c\pm} \text{---} \\ \text{---} \psi_{H^0} \text{---} \times \text{---} \tilde{z}^c \end{array} \Rightarrow -\frac{1}{16\pi^2} \frac{\epsilon^2}{(2n-1)^2} \int dx \frac{g}{c} (g_L + g_R) 4x(1-x) \quad (33)$$

3.2.4 Total neutral weak contribution

Again, as in case of QED, one can check that in the supersymmetric limit, discussed above, the contribution of the KK modes vanishes. Moreover, in the limit $g_L = -g_R = e$, the QED results are recovered.

Then, we sum up all the results of the neutral weak sector and obtain:

$$\Delta a_\mu^Z = \frac{g^2}{16\pi^2} (mR)^2 \frac{1}{c^2} \left(\frac{1}{4} - \frac{s^2}{3} + \frac{2s^4}{3} \right) \frac{\pi^2}{12} \quad (34)$$

3.3 Charged Weak sector

In this section we consider all the diagrams involving $W^\pm$ hypermultiplets fields and charged Higgs. The SM result [13] comes from graph *III*, where the loop contains a zero mode W boson and a neutrino and $g_+ = g/\sqrt{2}$, $g_- = 0$:

$$a_\mu^W = \frac{5g^2}{96\pi^2} \frac{m^2}{M_W^2} \quad (35)$$

3.3.1 KK vectors

The interaction terms involving the KK vectors are quite simple

$$\frac{g}{\sqrt{2}} W_n^\rho \mu_0 \sigma_\rho \bar{\psi}_{\nu,n} - \frac{g}{2\sqrt{2}} \frac{\epsilon}{2n} W_n^\rho \mu_0^c \sigma_\rho \bar{\psi}_{\nu,n} \quad (36)$$

and the external and internal cross diagrams are readily computed using graph *III*:

$$\begin{array}{c} \mu_0^c \text{---} \times \text{---} W_\rho^- \text{---} \mu_0 \\ \text{---} \psi_\nu \end{array} \Rightarrow -\frac{g^2}{16\pi^2} \frac{\epsilon^2}{(2n)^2} \int dx (1-x)^2 (x-3/2) \quad (37)$$

$$\begin{array}{c} \mu_0^c \text{---} W_\rho^- \text{---} \mu_0 \\ \text{---} \psi_\nu^c \times \text{---} \psi_\nu \end{array} \Rightarrow \frac{g^2}{16\pi^2} \frac{\epsilon^2}{(2n)^2} \int dx \frac{3}{2} (1-x)^2 \quad (38)$$

There also are diagrams of type V. The only nonvanishing one has a Higgs scalar, since the contributions of W_5 and σ_W cancel out:

$$\begin{array}{c} \mu_0^c \text{---} \phi_{H^+} \cdots \cdots W^- \text{---} \mu_0 \\ \text{---} \psi_\nu \end{array} \Rightarrow \frac{g^2}{16\pi^2} \frac{\epsilon^2}{(2n)^2} \int dx (1-x)^2 \quad (39)$$

3.3.2 Scalars

The gauge fixing procedure works as in the Z 's case: the interaction lagrangian terms only allow for an external cross contribution:

$$g(\psi_\nu^c \Sigma^+ \psi_L + \psi_L^c \Sigma^- \psi_\nu + h.c.)$$

$$\begin{array}{c} \mu_0^c \text{---} \times \text{---} \Sigma_w^+ \text{---} \mu_0 \\ \text{---} \psi_\nu^c \end{array} \Rightarrow -\frac{g^2}{16\pi^2} \frac{\epsilon^2}{(2n)^2} \int dx x(1-x)^2 \quad (40)$$

3.3.3 Fermions

The charginos $\tilde{w}^-$, $\tilde{w}_c^-$, ψ_{H^+} , $\psi_{H^+}^c$ mix for each n according to:

$$\mathcal{L}_{mass} = (\psi_{H^+}, \tilde{w}_c^-) \begin{pmatrix} \frac{2n-1}{R} & \sqrt{2}M_W \\ -\sqrt{2}M_W & \frac{2n-1}{R} \end{pmatrix} \begin{pmatrix} \psi_{H^+}^c \\ \tilde{w}^- \end{pmatrix} - \frac{2n-1}{R} \tilde{w}_c^+ \tilde{w}^+ \quad (41)$$

They enter the following interaction terms

$$\begin{aligned}
& g\phi_{\nu,n}^\dagger \tilde{w}_n^+ \mu_0 - g \frac{\epsilon}{2(2n-1)} \phi_{\nu,n}^\dagger \tilde{w}_n^- \mu_0^c + \\
& g\phi_{\nu,n}^c \tilde{w}_{c,n}^- \mu_0 + g \frac{\epsilon}{2(2n-1)} \phi_{\nu,n} \tilde{w}_{c,n}^+ \mu_0^c \\
& - \frac{2m}{v} \mu_0^c \psi_{H^+,n}^c \phi_{\nu,n}^{c\dagger}
\end{aligned} \tag{42}$$

The external cross diagrams are of the usual form

$$\left. \begin{array}{c} \text{Diagram 1: } \mu_0^c \text{ --- } \times \text{ --- } \tilde{w}^+ \text{ --- } \mu_0 \text{ with } \phi_\nu \text{ loop} \\ \text{Diagram 2: } \mu_0^c \text{ --- } \times \text{ --- } \tilde{w}_c^- \text{ --- } \mu_0 \text{ with } \phi_\nu^c \text{ loop} \end{array} \right\} \Rightarrow \frac{g^2}{16\pi^2} \frac{\epsilon^2}{(2n-1)^2} \int dx x(1-x)^2 \times 2 \tag{43}$$

and there are three internal cross diagrams with propagators $(\tilde{w}^+)^\dagger \text{---} \tilde{w}_c^+$, $(\tilde{w}_c^-)^\dagger \text{---} \tilde{w}^-$ and $(\tilde{w}_c^-)^\dagger \text{---} \phi_{H^+}^c$:

$$\begin{array}{c} \text{Diagram 3: } \mu_0^c \text{ --- } \tilde{w}_c^+ \text{ --- } \times \text{ --- } \tilde{w}^+ \text{ --- } \mu_0 \text{ with } \phi_\nu \text{ loop} \end{array} \Rightarrow \frac{g^2}{16\pi^2} \frac{\epsilon^2}{(2n-1)^2} \int dx (1-x)^2 \tag{44}$$

$$\begin{array}{c} \text{Diagram 4: } \mu_0^c \text{ --- } \tilde{w}^- \text{ --- } \times \text{ --- } \tilde{w}_c^- \text{ --- } \mu_0 \text{ with } \phi_\nu^c \text{ loop} \end{array} \Rightarrow -\frac{g^2}{16\pi^2} \frac{\epsilon^2}{(2n-1)^2} \int dx (1-x)^2 \tag{45}$$

$$\begin{array}{c} \text{Diagram 5: } \mu_0^c \text{ --- } \tilde{w}_c^- \text{ --- } \times \text{ --- } \psi_{H^+}^c \text{ --- } \mu_0 \text{ with } \phi_\nu^c \text{ loop} \end{array} \Rightarrow -\frac{g^2}{16\pi^2} \frac{\epsilon^2}{(2n-1)^2} \int dx 4(1-x)^2 \tag{46}$$

3.3.4 Total charged weak contribution

Again in this case, we correctly find that in the supersymmetric limit the sum of the KK states contributions vanishes.

We then gather all the contributions of the charged weak sector into the result:

$$\Delta a_\mu^W = -\frac{g^2}{16\pi^2}(mR)^2 \frac{7}{6} \frac{\pi^2}{12} \quad (47)$$

3.4 Result to leading order in $(M_W R)^2$

Collecting all partial results (20), (34) and (47) we have:

$$\Delta a_\mu^{KK} = -\frac{g^2}{192} \frac{m^2}{M_W^2} \frac{11 - 18s^2}{12c^2} (M_W R)^2 \quad (48)$$

Numerically, for $1/R = 370 \pm 70 \text{ GeV}$, it is

$$\Delta a_\mu^{KK} = 0.07^{+0.06}_{-0.02} \cdot a_\mu^{weak} = -(1.1^{+0.6}_{-0.3}) \cdot 10^{-10},$$

which means that, for any sensible value of R (see [14]), Δa_μ^{KK} is well inside the uncertainties that affect the SM prediction.

4 $M_W R$ corrections

In the previous section, we have neglected the ElectroWeak Symmetry Breaking in the masses of the KK modes for the gauge fields, since this was enough to get eq. (48). However, in this way we are neglecting contributions that in principle can be quite relevant, since the dimensionless expansion parameters $M_W R$ or $M_Z R$ can be as large as $1/3$. In this chapter we estimate these contributions. They are recovered if the full gauge fields masses are kept in the denominators of eq. (51)–(55), so the corrections to our previous results can be expressed in a simple way. Of course, these corrections only apply to the pure weak sector.

To begin with, let us focus on the neutral weak sector. First of all, what we have to do is to study the exact spectrum: defining the dimensionless parameter $\zeta = M_Z R$, for every n there is a vector boson Z_μ whose full mass squared is $(2n)^2 + \zeta^2$, in unit of $1/R^2$. The scalars fields masses are determined from eq. (26): the n^{th} KK mode of Σ_z has mass squared $(2n)^2 + \zeta^2$, while, for the real field φ_{H^0} , it has mass squared $(2n)^2 + \frac{3}{2}\zeta^2$. For the neutralinos, diagonalization of eq. (28) shows that there are two degenerate states with mass squared $(2n - 1)^2 + \zeta^2$. The rest of the computation proceeds as in the previous section. To get the modified results, it is sufficient to substitute the sum $\sum \frac{1}{(2n)^2}$ in eqs. (24), (25), (27), (30)–(33) with:

$$\sum \frac{1}{N^2 + \zeta^2} \quad \text{or} \quad \sum \frac{N^2}{(N^2 + \zeta^2)^2}$$

where $N = 2n$ for bosons and $N = 2n - 1$ for fermions. The first is the one to be used in all external cross diagrams, while for internal cross ones the formulae are a bit more involved. To be specific, in the case of the vectors, eq. (25) becomes:

$$-\frac{1}{16\pi^2}\epsilon^2 \int dx \left[\left(g_L g_R + \frac{1}{2}(g_L + g_R)^2 \right) 8x(1-x) \frac{1}{(2n)^2 + \zeta^2} - g_L g_R 16x(1-x)^2 \frac{(2n)^2}{((2n)^2 + \zeta^2)^2} \right] \quad (49)$$

With a bit of algebra, all contributions can be recast in a common form and the net result is:

$$\Delta a_\mu^Z = \frac{1}{16\pi^2} \frac{m^2}{M_W^2} (M_W R)^2 \{ (g_L^2 + g_R^2) f_1(\zeta) + g_L g_R f_2(\zeta) \} \quad (50)$$

where the functions f_1 and f_2 are defined in app. B. This replaces eq. (34), with all corrections to any order in $(M_W R)^2$ included. Numerically, for $1/R = 370 \pm 70 \text{ GeV}$, the result is only about 3% smaller than eq. (34). In the case of the charged weak sector, the calculation is very similar, so we expect the correction to be not so far from a few percent.

5 Conclusion

We have performed the calculation at one loop of the corrections to muon anomalous magnetic moment coming from the presence of one extra dimension in the model in ref. [1], to first order in $(mR)^2$ and in $(M_W R)^2$. We have also shown that the complete calculation in $(M_W R)^2$ yields small deviations, so to our purposes the simple analytic form given in eq. (48) is a good approximation. The corrections are relatively small, for any sensible value of R , at the 10% level of the pure weak contribution in the SM, and well inside the uncertainties of the hadronic contribution.

Appendix A

In this appendix we list the contributions to a_μ from the five graphs² in figure 1:

$$\Delta a_\mu^I = \frac{Q_s}{16\pi^2} m^2 \int_0^1 dx \left[\frac{h_+^2 + h_-^2}{2} 2x(1-x)^2 + \frac{m_f}{m} h_+ h_- 2x(1-x) \right] \frac{1}{xm_f^2 + (1-x)m_s^2 - x(1-x)m^2} \quad (51)$$

$$\Delta a_\mu^{II} = -\frac{Q_f}{16\pi^2} m^2 \int_0^1 dx \left[\frac{h_+^2 + h_-^2}{2} 2x(1-x)^2 + \frac{m_f}{m} h_+ h_- 2(1-x)^2 \right] \frac{1}{xm_s^2 + (1-x)m_f^2 - x(1-x)m^2} \quad (52)$$

²Similar results can also be found in refs. [15].

$$\Delta a_\mu^{III} = \frac{Q_v}{16\pi^2} m^2 \int_0^1 dx \left[\frac{g_+^2 + g_-^2}{2} 2(1-x)^2(2x-3) + g_+ g_- \frac{m_f}{m} 6(1-x)^2 \right] \frac{1}{xm_f^2 + (1-x)m_v^2 - x(1-x)m^2} \quad (53)$$

$$\Delta a_\mu^{IV} = -\frac{Q_f}{16\pi^2} m^2 \int_0^1 dx \left[\frac{g_+^2 + g_-^2}{2} (4x(1-x)^2 - 8x(1-x)) + \frac{m_f}{m} g_+ g_- 8x(1-x) \right] \frac{1}{xm_v^2 + (1-x)m_f^2 - x(1-x)m^2} \quad (54)$$

$$\Delta a_\mu^V = \frac{Q_v}{16\pi^2} m\tau \int_0^1 dx (g_+ h_+ + g_- h_-) (1-x)^2 \frac{1}{xm_f^2 + (1-x)m_v^2 - x(1-x)m^2} \quad (55)$$

where $Q_{s,f,v}$ are the electric charges of the particles emitting the zero-mode photon and $m_{s,f,v}$ are the masses in the loop. Note that in eq. (55) we assume $m_s = m_v$. The couplings are defined by:

$$\begin{aligned} & v_\rho \bar{f} \gamma^\rho [g_+ \mathcal{P}^+ + g_- \mathcal{P}^-] \mu \\ & s \bar{f} [h_+ \mathcal{P}^+ + h_- \mathcal{P}^-] \mu \\ & e \tau s^+ W_\rho^- A^\rho \\ & e [g^{\nu\rho} (k - p_+)^{\sigma} + g^{\rho\sigma} (p_+ - p_-)^{\nu} + g^{\sigma\nu} (p_- - k)^{\rho}] W_\rho^+ W_\sigma^- A_\nu \end{aligned}$$

where $P^\pm$ are projectors on the two Weyl components of the Dirac fields.

Appendix B

In this appendix we list the functions to be inserted in eq. (50):

$$\begin{aligned} f_1 &= \frac{1}{\zeta^8} \int_0^\zeta dy \, 2(\zeta^4 - y^4) \left(\pi y^2 \tanh\left(\frac{\pi}{2}y\right) - \pi y^2 \coth\left(\frac{\pi}{2}y\right) + 2y \right) = \\ &= \frac{4}{3\zeta^2} + \frac{1}{3\pi^6 \zeta^8} \left[48\pi^5 \zeta^5 \left(\text{Li}_2(-e^{-\pi\zeta}) - \text{Li}_2(e^{-\pi\zeta}) \right) + 336\pi^4 \zeta^4 \left(\text{Li}_3(-e^{-\pi\zeta}) - \text{Li}_3(e^{-\pi\zeta}) \right) \right. \\ &\quad + 1440\pi^3 \zeta^3 \left(\text{Li}_4(-e^{-\pi\zeta}) - \text{Li}_4(e^{-\pi\zeta}) \right) + 4320\pi^2 \zeta^2 \left(\text{Li}_5(-e^{-\pi\zeta}) - \text{Li}_5(e^{-\pi\zeta}) \right) + \\ &\quad + 8640\pi \zeta \left(\text{Li}_6(-e^{-\pi\zeta}) - \text{Li}_6(e^{-\pi\zeta}) \right) + 8640 \left(\text{Li}_7(-e^{-\pi\zeta}) - \text{Li}_7(e^{-\pi\zeta}) \right) + \\ &\quad \left. - 42\pi^4 \zeta^4 \zeta(3) + 17145 \zeta(7) \right] \end{aligned}$$

$$\begin{aligned}
f_2 &= \frac{1}{\zeta^8} \int_0^\zeta dy \, 8(y^2 - \zeta^2)(3y^2 - 2\zeta^2) \left(\pi y^2 \tanh\left(\frac{\pi}{2}y\right) - \pi y^2 \coth\left(\frac{\pi}{2}y\right) + 2y \right) = \\
&= \frac{4}{\zeta^2} - \frac{4}{\pi^6 \zeta^8} \left[8\pi^5 \zeta^5 \left(\text{Li}_2(-e^{-\pi\zeta}) - \text{Li}_2(e^{-\pi\zeta}) \right) + 136\pi^4 \zeta^4 \left(\text{Li}_3(-e^{-\pi\zeta}) - \text{Li}_3(e^{-\pi\zeta}) \right) + \right. \\
&\quad + 960\pi^3 \zeta^3 \left(\text{Li}_4(-e^{-\pi\zeta}) - \text{Li}_4(e^{-\pi\zeta}) \right) + 3840\pi^2 \zeta^2 \left(\text{Li}_5(-e^{-\pi\zeta}) - \text{Li}_5(e^{-\pi\zeta}) \right) + \\
&\quad + 8640\pi \zeta \left(\text{Li}_6(-e^{-\pi\zeta}) - \text{Li}_6(e^{-\pi\zeta}) \right) + 8640 \left(\text{Li}_7(-e^{-\pi\zeta}) - \text{Li}_7(e^{-\pi\zeta}) \right) + \\
&\quad \left. + 28\pi^4 \zeta^4 \zeta(3) - 930\pi^2 \zeta^2 \zeta(5) + 17145 \zeta(7) \right]
\end{aligned}$$

where $\zeta(n)$ is the Riemann zeta function, not to be confused here with our variable $\zeta = M_Z R$. In the integrals, the hyperbolic tangent is produced by the sum over the fermionic KK states, while the cotangent and the “ $2y$ ” pieces by the bosonic states.

Acknowledgements

We would like to thank Riccardo Barbieri and Riccardo Rattazzi for useful discussions and suggestions. We are also grateful to Gilberto Colangelo for informations about the present status of SM $(g-2)_\mu$. This work was supported by the EC under the RTN contract HPRN-CT-2000-00148.

References

- [1] R.Barbieri, L.J.Hall, Y.Nomura, *Phys.Rev.***D63**:105007,2001, arXiv:hep-ph/0011311;
- [2] R.Barbieri, G.Cacciapaglia, A.Romito, arXiv:hep-ph/0107148;
- [3] G.Cacciapaglia, M.Cirelli, G.Cristadoro, arXiv:hep-ph:0111287;
- [4] H.N.Brown et al. (Muon g-2 Collaboration), *Phys.Rev.Lett.* **86**:2227-2231,2001, arXiv:hep-ex/0102017;
A.Czarnecki, W.J.Marciano, *Phys.Rev.* **D64**:013014,2001, arXiv:hep-ph/0102122;
- [5] P.Nath, M.Yamaguchi, *Phys.Rev.* **D60**:116006,1999, arXiv:hep-ph/9903298;
R.Casadio, A.Gruppuso, G.Venturi, *Phys. Lett.* **B495**:378, 200, arXiv:hep-th/0010065;
Seong Chan Park, H.S.Song, *Phys. Lett.* **B506**:99,2001, arXiv:hep-ph/0103072;
K.Agashe, N.G.Deshpande, G.-H.Wu, *Phys.Lett.* **B511**:85,2001, arXiv:hep-ph/0103235;
T.Appelquist, B.A.Dobrescu, *Phys.Lett* **B516**:85,2001, arXiv:hep-ph/0106140;
- [6] J.Prades, invited talk at “Kaon 2001”, 6-13 June 2001, Pisa, Italy, arXiv:hep-ph/0108192;
- [7] for a recent review: W.J.Marciano, B.L.Roberts, arXiv:hep-ph/0105056;
- [8] M.Knecht, A.Nyffeler, arXiv:hep-ph/0111058;
M.Knecht, A.Nyffeler, M.Perrottet, E.de Rafael, arXiv:hep-ph/0111059;
- [9] I.Blokland, A.Czarnecki and K.Melnikov, arXiv:hep-ph/0112117;
- [10] C.A.Scrucca, M.Serone, L.Silvestrini, F.Zwirner, arXiv:hep-th/0110073;
- [11] J.Schwinger, *Phys.Rev.* **73**:416-417,1948;
Phys.Rev. **76**:790-817:1949;
- [12] G.Altarelli, N.Cabibbo, L.Maiani, *Phys.Lett.* **B40**:415,1972;
- [13] S.J.Brodsky, J.D.Sullivan, *Phys.Rev.* **156**:1644,1967;
- [14] R.Barbieri, L.J.Hall, Y.Nomura, arXiv:hep-ph/0110102;
- [15] I.Vendramin, *Nuovo Cim.* **A101**:731,1989;
K.R.Lynch, arXiv:hep-ph/0108080 and arXiv:hep-ph/0108081 .